

Selected reading on characterization of forage legumes

Dawson IK, Simons AJ, Waugh R, Powell W. 1995. Diversity and genetic differentiation among subpopulations of *Gliricidia sepium* revealed by PCR-based assays. *Heredity* 74: 10-18.

Edye LA, Burt RL, Nicholson CHL, Williams RJ, Williams WT. 1974. Classification of the *Stylosanthes* collection. 1928-69. CSIRO Aust. Div. Trop. Agron. Tech. Pap. No. 15, 1-28.

Gillies ACM, Abbott RJ. 1996. Phylogenetic relationships in the genus *Stylosanthes* (Leguminosae) based upon chloroplast DNA variation. *Pl. Syst. Evol.* 200: 193-211.

Hakiza JJ, Lazier JR, Sayers AR. 1988a. Characterization and evaluation of forage legumes in Ethiopia: Preliminary examination of variation between accessions of *Stylosanthes fruticosa* (Retz.) Alston. Pages 174-191 in B.H. Dzewela, ed. African Forage Plant Genetic Resources, Evaluation of Forage Germplasm and Extensive Livestock Production Systems. ILCA, Addis Ababa, Ethiopia.

Hakiza JJ, Lazier JR, Sayers AR. 1988b. Characterization and preliminary evaluation of accessions of *Zornia* species from the ILCA collection. Pages 149-173 in B.H. Dzewela, ed. African Forage Plant Genetic Resources, Evaluation of Forage Germplasm and Extensive Livestock Production Systems. ILCA, Addis Ababa, Ethiopia.

Heering H, Nokoe S, Mohammed J. 1996a. The classification of a *Sesbania sesban* collection. 1. Morphological attributes and their taxonomic significance. *Trop. Grassl.* 30: 206-214.

Heering H, Reed JD, Hanson J. 1996b. Differences in *Sesbania sesban* accessions in relation to their phenolic concentration and HPLC fingerprints. *J. Sci. Food Agric.* 71: 92-98.

IBPGR. 1992. Descriptors for White Clover (*Trifolium repens*). International Board for Plant Genetic Resources, Rome, Italy. 52 pp.

Jones PG, Rebgetz R, Maass BL, Kerridge PC. 1996. Genetic diversity in *Stylosanthes* species: a GIS mapping approach. Paper presented at the First International Symposium on Tropical Savannas, Brasília, DF, Brazil, 24-29 March 1996.

Kahurananga J, Tsehay A. 1991. Variation in flowering time, dry matter and seed yield among annual *Trifolium* species, Ethiopia. *Trop. Grasslands* 25, 20-25.

Kaufman L, Rousseeuw PJ. 1990. Finding Groups in Data: An Introduction to Cluster Analysis. John Wiley & Sons, New York, USA. 342 pp.

Kazan K, Manners JM, Cameron DF. 1993a. Genetic variation in agronomically important species of *Stylosanthes* determined using amplified polymorphic DNA markers. *Theor. Appl. Genet.* 85: 882-888.

Kazan K, Manners JM, Cameron DF. 1993b. Genetic relationships and variation in the *Stylosanthes guianensis* (Aubl.) Sw. species complex assessed by random amplified polymorphic DNA. *Genome* 36: 43-49.

Larbi A, Ochang J, Hanson J, Lazier J. 1992. Agronomic evaluation of *Neonotonia wightii*, *Stylosanthes scabra* and *S. guianensis* accessions in Ethiopia. *Trop. Grasslands* 26, 115-119.

Larbi A, Thomas D, Hanson J. 1993b. Forage potential of *Erythrina abyssinica*: intake, digestibility and growth rates for stall-fed sheep and goats in southern Ethiopia. *Agroforestry Systems* 21, 263-270.

- Larbi A, Thomas D, Ochang J. 1993a. Agronomic evaluations of *Chamaecrista rotundifolia* accessions under grazing in southern Ethiopia. *E. Afr. agric. For. J.* 59, 75-78.
- Lowe AJ, Hanotte O, Guarino L. 1996. Standardisation of molecular genetic techniques for the characterisation of germplasm collections: The case for Random Amplified Polymorphic DNA (RAPD). *Plant Genetic Resources Newsletter*, 107: 50-54.
- Luckow, M. (1993) Monograph of *Desmanthus* (Leguminosae-Mimosoideae). *Systematic Botany Monographs* 38. 166pp.
- Maass BL. 1989. Die tropische Weideleguminose *Stylosanthes scabra* Vog. - Variabilität, Leistungsstand und Möglichkeiten züchterischer Verbesserung. Sonderheft 97. *Landbau-forschung Völknerode, Braunschweig, Germany.* 189 p.
- Maass BL, Ocampo CH. 1995. Isozyme polymorphism provides fingerprints for germplasm of *Arachis glabrata* Benth. *Genet. Resour. Crop Evol.* 42: 77-82.
- Maass BL, Jamnadass RH, Hanson J, Pengelly BC. 2005. Determining sources of diversity in cultivated and wild *Lablab purpureus* related to provenance of germplasm by using amplified fragment length polymorphism. *Genetic Resources and Crop Evolution*, 52: 683–695.
- Maass BL, Schultze-Kraft R. 1993. Characterisation and preliminary evaluation of a large germplasm collection of the tropical forage legume *Stylosanthes scabra* Vog. *Proc. XVII Int. Grass. Cong., New Zealand and Australia*, pp 2151-2153.
- Maass BL, Torres AM, Ocampo CH. 1993. Morphological and isozyme characterisation of *Arachis pintoii* Krap. *et Greg. nom. nud.* germplasm. *Euphytica* 70: 43-52.
- Macqueen DJ. 1993. *Calliandra* Series Racemosae: Taxonomic information; OFI seed collections; trial design. Oxford Forestry Institute, Oxford. (Unpublished Report).
- McDonald MB, Elliot LJ, Sweeney PM. 1994. DNA extraction from dry seeds for RAPD analyses in varietal identification studies. *Seed Sci. & Technol.* 22, 171-176.
- Pengelly BC, Eagles DA. 1995. Geographical distribution and diversity in a collection of the tropical legume *Macroptilium gracile* (Poeppig ex Benth) Urban. *Aust. J. Agric. Res.* 46: 569-80.
- Pengelly BC, Maass BL. 2001. *Lablab purpureus* (L.) Sweet – diversity, potential use and determination of a core collection of this multi-purpose tropical legume. *Genetic Resources and Crop Evolution* 48:261-272.
- Peters M, Tarawali SA, Alkamper J. 1994a. Evaluation of tropical pasture legumes for fodder banks in subhumid Nigeria. 1. Accessions of *Centrosema brasilianum*, *C. pascuorum*, *Chamaecrista rotundifolia* and *Stylosanthes hamata*. *Trop. Grasslands* 28, 65-73.
- Peters M, Tarawali SA, Alkamper J. 1994b. Evaluation of tropical pasture legumes for fodder banks in subhumid Nigeria. 2. Accessions of *Aeschynomene hystrix*, *Centrosema acutifolium*, *C. pascuorum*, *Stylosanthes guianensis* and *S. hamata*. *Trop. Grasslands* 28, 74-79.
- Pizarro EA, Coradin L. (eds.). 1996. Potencial del género *Cratylia* como leguminosa forrajera. Proceedings of a workshop 17-20 July 1995, Brasilia, D.F., Brazil. Working document. CIAT, Cali, Colombia. 118 p.
- Plumb VE, Wood CD, Gay C, Hanson J, Heering JH. 1996. The use of HPLC-derived phenolic profiles as a means of classifying *Sesbania* accessions. *J. Sci. Food Agric.* 70: 481-486.

Reed JD, Krueger C, Rodriguez G, Hanson J. 2000. Secondary plant compounds and forage evaluation. In: Givens, D.I., Owen, E., Axford, R.F.E. and Omed, H.M. (eds). Forage Evaluation in Ruminant Nutrition. CABI Publishing, UK. pp 433-448.

Schultze-Kraft R, Belalcázar J. 1988. Germplasm collection and preliminary evaluation of the pasture legume *Centrosema brasilianum* (L.) Benth. Tropic. Agric. (Trinidad) 65(2): 137-144.

Schultze-Kraft R, Benavides G. 1988. Germplasm collection and preliminary evaluation of *Desmodium ovalifolium* Wall. Genet. Resour. Commun. 12: 1-20.

Tarawali SA. 1991. Preliminary agronomic evaluation of forage legumes for subhumid West Africa. Trop. Agric. 68, 88-94.

Torres AM, Belalcázar J, Andrade M, Ortiz A. 1995. Variabilidad del germoplasma del género *Chamaecrista* (Breyne) Moench (Leguminosae: Caesalpinioideae) conservado en el CIAT. Pages 403-407 in Memorias del I Congreso Nacional sobre Biodiversidad, Cali, Colombia, 4-8 December 1994.

Van de Wouw M, Maxted N, Ford-Lloyd BV. 2003. [Agro-morphological characterisation of common vetch and its close relatives](#). *Euphytica* 130: 281-292.