

Concentration, temperatures and period of staining with tetrazolium solution (for Annex 1 crops of the International Treaty on PGRFA).

<i>Crop</i>	<i>Species</i>	<i>Preconditioning</i>	<i>Staining</i>
Barley	<i>Hordeum vulgare</i>	Imbibe or soak, 6-18h	0.5%, 3h, 30°C
Beans	<i>Phaseolus</i> spp.	Imbibe 18-24h, then soak, 2-3h	0.5-1%, 6-24h, 30°C
Brassica	<i>Brassica</i> spp.	Imbibe or soak, 16- 18h	0.5-1%, 3-6h, 30°C
Chickpea	<i>Cicer arietinum</i>	Imbibe or soak, 18h	1%, 6-24h, 30°C
Cowpea	<i>Vigna unguiculata</i>	Soak, 22h	0.5-1%, 16-24h, 30°C
Eggplant	<i>Solanum melongena</i>	Imbibe or soak, 18h	0.5-1%, 6-24h, 30°C
Faba bean	<i>Vicia faba</i>	Soak, 22h	0.5-1%, 16-24h, 30°C
Finger millet	<i>Eleusine corocana</i>	Soak, 18h, 5°C	0.5%, 3h, 30°C
Lentil	<i>Lens culinaris</i>	Imbibe, 18h, then soak, 2-3h	1%, 6-24h, 30°C
Maize	<i>Zea mays</i>	Imbibe or soak, 18h	0.5-1%, 2-6h, 30°C
Pea	<i>Pisum sativum</i>	Imbibe 18-24h, then soak, 2-3h	0.5-1%, 6-24h, 30°C
Pearl millet	<i>Pennisetum glaucum</i>	Imbibe or soak, 6-18h	0.5-1%, 6-24h, 30°C
Rice	<i>Oryza sativa</i>	Imbibe or soak, 18h	0.5%, 3h, 30°C
Rye	<i>Secale cereale</i>	Imbibe or soak, 6-18h	0.5%, 2-3h, 30°C
Sorghum	<i>Sorghum bicolor</i>	Imbibe, 16h, 30°C	0.5-1%, 0.5-1h, 40°C
Sugar beet	<i>Beta vulgaris</i>	Imbibe or soak, 16- 18h	1%, 24-48h, 30°C
Sunflower	<i>Helianthus annuus</i>	Imbibe or soak, 18h	0.5-1%, 3-6, 30°C
Triticale	<i>Triticosecale</i>	Imbibe or soak, 6-18h	0.5%, 2-4h, 30°C
Wheat	<i>Triticum aestivum</i>	Imbibe or soak, 6-18h	0.5%, 2-4h, 30°C