

Suggested method of moisture determination for important crop and forage species (ISTA 2005)

<i>Low-constant temperature oven method</i>		
Brassicas Castor (<i>Ricinus</i>)* Pepper (<i>Capsicum</i>) Cotton (<i>Gossypium</i>)* Eggplant (<i>Solanum</i>)	Falseflax (<i>Camelina</i>) Flax (<i>Linum</i>) Groundnut (<i>Arachis</i>)* Onion (<i>Allium</i>) Radish (<i>Raphanus</i>)	Sesame (<i>Sesamum</i>) Soyabean (<i>Glycine</i>)* All tree species
<i>High-constant temperature oven method</i>		
Alfalfa (<i>Medicago</i>) Asparagus (<i>Asparagus</i>) Barley (<i>Hordeum</i>)* Bean (<i>Phaseolus</i>)* Beet (<i>Beta</i>) Bentgrass (<i>Agrostis</i>) Bermuda grass (<i>Cynodon</i>) Black salsify (<i>Scorzonera</i>) Bluegrass (<i>Poa</i>) Brome (<i>Bromus</i>) Buckwheat (<i>Fagopyrum</i>)* Canarygrass (<i>Phalaris</i>) Caraway (<i>Carum</i>) Carrot (<i>Daucus</i>) Chervil (<i>Anthriscus</i>) Chickory (<i>Cichorium</i>) Chickpea (<i>Cicer</i>)* Clover (<i>Trifolium</i>)	Cocksfoot (<i>Dactylis</i>) Cress (<i>Lepidium</i>) Crested dogtail (<i>Cynosurus</i>) Cucumber (<i>Cucumis</i>) Cumin (<i>Cuminum</i>) Dallisgrass (<i>Paspalum</i>) Fescue (<i>Festuca</i>) Foxtail (<i>Alopecurus</i>) Lettuce (<i>Lactuca</i>) Lupin (<i>Lupinus</i>)* Maize (<i>Zea</i>)* Millet (<i>Panicum</i>) Oat (<i>Avena</i>)* Parsley (<i>Petroselinum</i>) Pea (<i>Pisum</i>)* Rhodes grass (<i>Chloris</i>) Rice (<i>Oryza</i>)*	Rye (<i>Secale</i>)* Ryegrass (<i>Lolium</i>) Sainfoin (<i>Onobrychis</i>) Serradella (<i>Ornithopus</i>) Sorghum (<i>Sorghum</i>)* Squash (<i>Cucurbita</i>) Sweetclover (<i>Melilotus</i>) Tall oatgrass (<i>Arrhenatherum</i>) Timothy grass (<i>Phleum</i>) Tomato (<i>Lycopersicon</i>) Trefoil (<i>Lotus</i>) Tufted hairgrass (<i>Deschampsia</i>) Velvetgrass (<i>Holcus</i>) Vetch (<i>Vicia</i>)* Watermelon (<i>Citrullus</i>)* Wheat (<i>Triticum</i>)*

*grinding required